

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/52926>

Please be advised that this information was generated on 2017-12-06 and may be subject to change.

ROBERT P. BAUGHMAN
University of Cincinnati Medical Center
Cincinnati, Ohio

References

1. Baughman RP, Drent M, Kavuru M, Judson MA, Costabel U, du Bois R, Albera C, Brutsche M, Davis G, Donohue JF, *et al*. Infliximab therapy in patients with chronic sarcoidosis and pulmonary involvement. *Am J Respir Crit Care Med* 2006;174:795–802.
2. Pietinalho A, Tukiainen P, Haahtela T, Persson T, Selroos O; Finnish Pulmonary Sarcoidosis Study Group. Early treatment of stage II sarcoidosis improves 5-year pulmonary function. *Chest* 2002;121:24–31.
3. Gibson GJ, Prescott RJ, Muers MF, Middleton WG, Mitchell DN, Connolly CK, Harrison BD. British Thoracic Society Sarcoidosis study: effects of long term corticosteroid treatment. *Thorax* 1996;51:238–247.
4. Muers MF, Middleton WG, Gibson GJ, Prescott RJ, Mitchell DN, Connolly CK, Harrison BD. A simple radiographic scoring method for monitoring pulmonary sarcoidosis: relations between radiographic scores, dyspnoea grade and respiratory function in the British Thoracic Society Study of Long-Term Corticosteroid Treatment. *Sarcoidosis Vasc Diffuse Lung Dis* 1997;14:46–56.
5. Drent M, Costabel U, Shipley R, Desai S, Judson M, du Bois R, Lo KH, Watt R, Baughman RP. Changes of chest roentgenogram during a double blind randomized trial of infliximab for chronic pulmonary sarcoidosis. *Eur Respir J* 2006;28:111S.
6. Baughman RP, Lower EE. Infliximab for refractory sarcoidosis. *Sarcoidosis Vasc Diffuse Lung Dis* 2001;18:70–74.
7. O'Shea FD, Marras TK, Inman RD. Pulmonary sarcoidosis developing during infliximab therapy. *Arthritis Rheum* 2006;55:978–981.

Impact of New American Thoracic Society Diagnostic Criteria on Management of Nontuberculous Mycobacterial Infection

To the Editor:

The American Thoracic Society has recently published new diagnostic criteria for nontuberculous mycobacterial (NTM) diseases (1). We retrospectively assessed the impact of the new criteria on clinical practice by reviewing medical files of all patients in whom *Mycobacterium chelonae*, *M. abscessus*, *M. simiae*, or *M. szulgai* were isolated between January 1999 and January 2005 in The Netherlands. Isolates were earlier identified at the National Mycobacteria Reference Laboratory (RIVM), by either reverse line blot assay (InnoLipa Mycobacteria v2; Innogenetics, Ghent, Belgium) or 16S gene sequencing, or by a self-sufficient hospital laboratory. Approximately 7.7% (9/117) more pulmonary patients met the 2007 diagnostic criteria (1) than the 1997 criteria (2), although this differed by NTM species (Table 1).

TABLE 1. FULFILLMENT OF THE DIAGNOSTIC CRITERIA, BY SPECIES AND SITE

Species	Site	n	Age (yr)	1997 ATS+, n (%)	2007 ATS+, n (%)
<i>Mycobacterium chelonae</i>	P	35	57	5 (14)	7 (20)
	EP	11	51	8 (73)	7 (64)
<i>M. abscessus</i>	P	39	51	11 (28)	13 (33)
	EP	10	52	9 (90)	9 (90)
<i>M. szulgai</i>	P	15	60	8 (53)	11 (73)
	EP	6	44	6 (100)	5 (83)
<i>M. simiae</i>	P	28	65	6 (21)	8 (29)

Definition of abbreviations: Age = mean age; ATS+ = American Thoracic Society diagnostic criteria met; EP = extrapulmonary isolates; P = pulmonary isolates.

The new diagnostic criteria contain more specific radiological criteria, whereas bacteriological criteria have become more lenient: a single NTM culture from bronchial washing fluid, in a well-defined class of patients, or two positive sputum cultures now suffice to establish the diagnosis. As more patients are likely to meet the new criteria, more patients might receive antimycobacterial treatment. The authors, however, repeatedly state that meeting these criteria does not, *per se*, necessitate the institution of therapy (1). This was not emphasized in the former statement. We believe such comments only add to the confusion for the clinician. Which factors should influence this treatment decision in individual patients? Ten years of experience with the former criteria apparently have not provided the answer to that question. More lenient criteria even increase the responsibility of physicians in the treatment decision.

Increased exposure to antimycobacterial drugs can harm patients, both in terms of adverse events and cost (3). The variable cure rates achievable per type of NTM disease and persistence of the conditions predisposing to NTM infection further complicate the assessment of the benefit of treatment. Centralization of expertise and easily accessible expert consultation are important, as acknowledged in the new statement. For extrapulmonary NTM isolates, no criteria for additional proof of infection are mentioned in the summarized criteria, contrary to the prior statement (2). This complicates assessment of the significance of NTM isolated from pleural fluid, stool, or gastric juice, without a context of disseminated disease, and in the HIV-negative patient, without histological proof of infection (Table 1). More research is needed to improve diagnosis and treatment of this increasing problem in clinical practice.

Conflict of Interest Statement: None of the authors has a financial relationship with a commercial entity that has an interest in the subject of this manuscript.

JAKKO VAN INGEN
Radboud University
Nijmegen Medical Centre
Nijmegen, The Netherlands
and
National Mycobacteria Reference Laboratory
Bilthoven, The Netherlands
MARTIN J. BOEREE
WIEL C. M. DE LANGE
P. N. RICHARD DEKHUIJZEN
Radboud University
Nijmegen Medical Centre
Nijmegen, The Netherlands
DICK VAN SOOLINGEN
National Mycobacteria Reference Laboratory
Bilthoven, The Netherlands

References

1. Griffith DE, Aksamit T, Brown-Elliott BA, Catanzaro A, Daley C, Gordin F, Holland SM, Horsburgh R, Huitt G, Iademaro MF, *et al*. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med* 2007;175:367–416.
2. American Thoracic Society. Diagnosis and treatment of disease caused by nontuberculous mycobacteria. *Am J Respir Crit Care Med* 1997;156:S1–S25.
3. van Crevel R, de Lange WCM, Vanderpuye NA, van Soolingen D, Hoogkamp-Korstanje JAA, van Deuren KM, Kullberg BJ, van Herwaarden C, van der Meer JW. The impact of nontuberculous mycobacteria on management of presumed pulmonary tuberculosis. *Infection* 2001;29:59–63.